

Merck Animal Health

Equine Products Reference Guide



Vaccines

Equi-Nile™ WNV WNV - KILLED

New!



Equi-Nile™ WNV with Havlogen®*

WEST NILE VIRUS VACCINE
KILLED FLAVIVIRUS CHIMERA

For vaccination of healthy horses 6 months of age or older; as an aid in reduction of disease, encephalitis, and viremia caused by West Nile Virus. Duration of immunity is under development. Foals nursing immune-dams should be vaccinated when maternal antibody levels will allow active immunization.

1 x 10 mL, 10 x 1 mL

EquiRab™ RABIES



1 mL dose

EquiRab™ with Havlogen®*

RABIES VACCINE
KILLED VIRUS

For vaccination of healthy horses, 4 months of age or older, as an aid in the prevention of disease due to rabies virus up to 14 months following vaccination.

1 x 10 mL, 10 x 1 mL

Super-Tet® TETANUS TOXOID



Super-Tet® with Havlogen®*

TETANUS TOXOID

Purified Tetanus Toxoid for vaccination of healthy horses, cattle, swine and sheep, 6 months of age or older, against Tetanus.

10 x 1 mL

Prestige® V + WNV EEE + WEE + TETANUS + EIV + EHV - 1&4 + WNV

New!



Prestige® V + WNV with Havlogen®*

ENCEPHALOMYELITIS - RHINOPNEUMONITIS -
INFLUENZA - WEST NILE VIRUS VACCINE,
EASTERN & WESTERN, KILLED FLAVIVIRUS CHIMERA,
TETANUS TOXOID

For vaccination of healthy horses 6 months of age or older; as an aid in the prevention of disease caused by Eastern and Western Encephalomyelitis viruses and Tetanus; as an aid in the control of respiratory disease caused by EIV, EHV-1 and EHV-4; as an aid in reduction of virus shedding of EIV, EHV-1 and EHV-4; and as an aid in reduction of disease, encephalitis, and viremia caused by West Nile Virus. DOI shown to be 6 months for Equine Influenza Virus.

1 x 10 mL, 10 x 1 mL

Prestige® V + VEE EEE + WEE + TETANUS + EIV + EHV - 1&4 + VEE



Prestige® V + VEE with Havlogen®*

ENCEPHALOMYELITIS - RHINOPNEUMONITIS -
INFLUENZA VACCINE
EASTERN, WESTERN, AND VENEZUELAN, KILLED VIRUS
TETANUS TOXOID

For vaccination of healthy horses, 6 months of age or older, as an aid in the prevention of Eastern, Western and Venezuelan Encephalomyelitis and Tetanus, and as an aid in the control of respiratory disease caused by Equine Influenza Virus and Equine Herpesvirus and as an aid in the reduction of EHV-1 and EHV-4 virus shedding and Influenza Virus shedding. DOI shown to be 6 months for Equine Influenza Virus.

1 x 10 mL, 10 x 1 mL

Prestige® V EEE + WEE + TETANUS + EIV + EHV - 1&4



Prestige® V with Havlogen®*

ENCEPHALOMYELITIS - RHINOPNEUMONITIS -
INFLUENZA VACCINE
EASTERN AND WESTERN, KILLED VIRUS
TETANUS TOXOID

For vaccination of healthy horses, 6 months of age or older, as an aid in the prevention of Eastern and Western Encephalomyelitis and Tetanus, as an aid in the control of respiratory disease caused by Equine Influenza Virus and Equine Herpesvirus and as an aid in the reduction of EHV-1 and EHV-4 virus shedding and Influenza Virus shedding. DOI shown to be 6 months for Equine Influenza Virus.

1 x 10 mL, 10 x 1 mL

Prestige® IV FoalShot EEE + WEE + TETANUS + EHV - 1&4



Prestige® IV Foalshot with Havlogen®*

ENCEPHALOMYELITIS - RHINOPNEUMONITIS
VACCINE
EASTERN AND WESTERN, KILLED VIRUS
TETANUS TOXOID

For vaccination of healthy horses, 6 months of age or older, as an aid in the prevention of disease caused by Eastern and Western Equine Encephalomyelitis Viruses and Tetanus, and as an aid in the control of respiratory disease caused by EHV-1 and EHV-4, and as an aid in reducing EHV-1 and EHV-4 virus shedding. Foals nursing immune-dams should be vaccinated when maternal antibody levels will allow active immunization.

1 x 10 mL

Prestige® II EIV + EHV - 1&4



Prestige® II with Havlogen®*

EQUINE RHINOPNEUMONITIS -
INFLUENZA VACCINE
KILLED VIRUS

For vaccination of healthy horses, 6 months of age or older, as an aid in the control of respiratory disease caused by Equine Influenza Virus and Equine Herpesvirus and as an aid in the reduction of EHV-1 and EHV-4 virus shedding and Influenza Virus shedding. DOI shown to be 6 months for Equine Influenza Virus.

1 x 10 mL, 10 x 1 mL

Prestige® EHV - 1&4



Prestige® with Havlogen®*

EQUINE RHINOPNEUMONITIS VACCINE
KILLED VIRUS

For vaccination of healthy horses, 6 months of age or older, as an aid in the control of respiratory disease caused by EHV-1 and EHV-4 and as an aid in the reduction of EHV-1 and EHV-4 virus shedding.

1 x 10 mL

Encevac® T + WNV
EEE + WEE + TETANUS + WNV

New!



Encevac® T + WNV with Havlogen®*

ENCEPHALOMYELITIS -
WEST NILE VIRUS VACCINE,

EASTERN & WESTERN, KILLED VIRUS, KILLED FLAVIVIRUS CHIMERA,
TETANUS TOXOID

For vaccination of healthy horses 6 months of age or older; as an aid in the prevention of disease caused by Eastern and Western Encephalomyelitis Viruses and Tetanus; and as an aid in reduction of disease, encephalitis, and viremia caused by West Nile Virus. Duration of immunity for the West Nile Virus fraction is under development. Foals nursing immune-dams should be vaccinated when maternal antibody levels will allow active immunization.

1 x 10 mL

Encevac® + WNV
EEE + WEE + WNV

New!



Encevac® + WNV with Havlogen®*

ENCEPHALOMYELITIS -
WEST NILE VIRUS VACCINE,

EASTERN & WESTERN, KILLED VIRUS, KILLED FLAVIVIRUS CHIMERA

For vaccination of healthy horses 6 months of age or older; as an aid in the prevention of disease caused by Eastern and Western Encephalomyelitis Viruses; and as an aid in reduction of disease, encephalitis, and viremia caused by West Nile Virus. Duration of immunity for the West Nile Virus fraction is under development. Foals nursing immune-dams should be vaccinated when maternal antibody levels will allow active immunization.

1 x 10 mL

Encevac®-T
EEE + WEE + TETANUS



Encevac®-T with Havlogen®*

ENCEPHALOMYELITIS VACCINE
EASTERN AND WESTERN, KILLED VIRUS

TETANUS TOXOID

For vaccination of healthy horses, 6 months of age or older, against Eastern and Western Equine Encephalomyelitis and Tetanus.

1 x 10 mL, 10 x 1 mL

Encevac® TC-4
EEE + WEE + TETANUS + EIV



Encevac® TC-4 with Havlogen®*

ENCEPHALOMYELITIS - INFLUENZA VACCINE

EASTERN AND WESTERN, KILLED VIRUS
TETANUS TOXOID

For vaccination of healthy horses, 6 months of age or older, as an aid in the prevention of Eastern and Western Equine Encephalomyelitis and Tetanus, and as an aid in the reduction of virus shedding and control of respiratory disease caused by Equine Influenza Virus. DOI shown to be 6 months for Equine Influenza Virus.

1 x 10 mL, 10 x 1 mL

Encevac® TC-4 + VEE
EEE + WEE + TETANUS + EIV + VEE



**Encevac® TC-4 + VEE
with Havlogen®***

ENCEPHALOMYELITIS - INFLUENZA VACCINE

EASTERN, WESTERN, AND VENEZUELAN, KILLED VIRUS
TETANUS TOXOID

For vaccination of healthy horses, 6 months of age or older, as an aid in the prevention of Eastern, Western and Venezuelan Equine Encephalomyelitis and Tetanus, and as an aid in reduction of virus shedding and the control of respiratory disease caused by Equine Influenza Virus. DOI shown to be 6 months for Equine Influenza Virus.

1 x 10 mL, 10 x 1 mL

Encevac® T + VEE
EEE + WEE + TETANUS + VEE



Encevac® T + VEE with Havlogen®*

ENCEPHALOMYELITIS VACCINE

EASTERN, WESTERN AND VENEZUELAN, KILLED VIRUS
TETANUS TOXOID

For vaccination of healthy horses, 6 months of age or older, against Eastern, Western and Venezuelan Equine Encephalomyelitis and Tetanus.

1 x 10 mL

Prodigy®
EHV - 1



Prodigy® with Havlogen®*

EQUINE RHINOPNEUMONITIS VACCINE

KILLED VIRUS

An inactivated, purified, concentrated, adjuvanted, tissue culture origin Equine Herpesvirus 1 (EHV-1) vaccine. For vaccination of healthy horses 6 months of age or older, as an aid in the prevention of abortion and respiratory disease associated with EHV-1 infection.

1 x 20 mL, 10 x 2 mL

Flu Avert®
EIV



Flu Avert® I.N. vaccine
EQUINE INFLUENZA VACCINE

Modified Live Virus - For Intranasal Use Only

For the vaccination of healthy horses, 11 months of age or older, as an aid in the reduction of clinical disease and viral shedding caused by equine influenza viruses of both the American and Eurasian lineages.

10 x 1 mL

*ADJUVANT- Intervet's Proprietary Technology

Pharmaceuticals

Panacur® POWERPAC / Paste 10%



Panacur® (fenbendazole) Powerpac Equine Dewormer

For the control of large strongyles (*Strongylus edentatus*, *S. equinus*, *S. vulgaris*), encysted early third stage (*hyobiotic*), late third stage and fourth stage cyathostome larvae, small strongyles, pinworms (*Oxyuris equi*), ascarids (*Parascaris equorum*), and arteritis caused by fourth stage larvae of *Strongylus vulgaris* in horses.

5 x 57 gm syringes

Paste 10% Equine Dewormer

For the control of large strongyles (*Strongylus edentatus*, *S. equinus*, *S. vulgaris*), encysted early third stage (*hyobiotic*), late third stage and fourth stage cyathostome larvae, small strongyles, pinworms (*Oxyuris equi*), ascarids (*Parascaris equorum*), and arteritis caused by fourth stage larvae of *Strongylus vulgaris* in horses.

12 x 25 gm, 12 x 92 gm syringes

Banamine® Paste/Injectable (flunixin meglumine)



Banamine® Paste/Injectable (flunixin meglumine)

For the alleviation of inflammation and pain associated with musculoskeletal disorders in the horse.

1 x 100 mL 794, 1 x 250 mL 794 injectables 30 gm syringe

Phenylzone® Paste (phenylbutazone)



Phenylzone® Paste (phenylbutazone)

For the relief of inflammatory conditions associated with the musculoskeletal system in horses.

1 x 12 gm syringe

E-SE®



E-SE® (SELENIUM, VITAMIN E)

For the control of the following clinical signs when associated with myositis (Selenium-Tocopherol Deficiency) syndrome: rapid respiration, profuse sweating, muscle spasms and stiffness, elevated SGOT.

1 x 100 mL 794

Dolorex®



Dolorex® (Butorphanol Tartrate)

A totally synthetic, centrally acting, narcotic agonist-antagonist analgesic with potent antitussive activity.

10 mg/mL 50 mL vial

Regu-Mate®



Regu-Mate®

(altrenogest)

Solution 0.22%

Is indicated to suppress estrus in mares.

1000 mL

Salix®



Salix® (furosemide)

Salix® is an effective diuretic possessing a wide therapeutic range. Pharmacologically it promotes the rapid removal of abnormally retained extracellular fluids. The rationale for the efficacious use of diuretic therapy is determined by the clinical pathology producing the edema. Salix® is indicated for the treatment of edema, (pulmonary congestion, ascites) associated with cardiac insufficiency and acute noninflammatory tissue edema.

The continued use of heart stimulants, such as digitalis or its glycosides is indicated in cases of edema involving cardiac insufficiency.

50 mL vial

Equine Technical Services

Craig Barnett, D.V.M.
Cynthia MacKenzie, D.V.M.
Joseph Manning, D.V.M., MBA
Wendy Vaala, V.M.D., Dipl. ACVIM
Christina Taylor, B.S., RVT

Veterinary Technical Service
866-349-3497

Merck Animal Health
556 Morris Ave., Summit, NJ 07901
Phone: 800-521-5767 www.merck-animal-health-usa.com

Merck Animal Health is a subsidiary of Merck & Co., Inc., Whitehouse Station, NJ, USA.
Except where noted, MERCK and all trademarks are registered trademarks or trademarks of Merck Sharp & Dohme Corp., a subsidiary of Merck & Co., Inc., Whitehouse Station, NJ, USA.
Avert is a registered trademark of Heska Corporation

*Adjuvant —Merck's Proprietary Technology

© 2011 Intervet International B.V, subsidiaries of Merck & Co., Inc., Whitehouse Station, NJ, USA. All rights reserved.
Part No. EQ-BIO-1083-F5, MAC 06/11

Marketing Team

Brett Whitehead
Director Equine and
Ag Retail Business
Joe Ferraro
Equine Product Manager

Customer Service
800-521-5767

Equine Sales Team

Ron McDaniel
National Sales Manager
— Equine

Jeff Baldrige
308-530-0407
NE, ND, SD, MO, MN, IA, KS

Kip Boule
407-448-2722
FL, GA, AL

Booth Farwell
443-686-1014
N. VA, DE, MD, NJ, E. PA, WV

Pete Hale
512-917-1998
TX, NM

Eric Johnson
719-660-8018
CO, UT, WY, MT

Trina Lally
302-521-9414
NY, New England

Allison McCosker
530-520-0472
N. CA, N. NV, OR, WA, ID

Steve Montemarano
740-972-6554
W. PA, OH, IN, IL, MI, WI

Gail Murray BS, RVT
859-608-8901
KY, TN

Jessica Steffien
909-227-0023
S. CA, S. NV, AZ, HI

Andrew Stetson
919-448-7039
S. VA, NC, SC

Mark Williams
214-384-4524
E. TX, MS, LA, AR, OK

Inside Sales Team

Amy Lenhart
Office: 800-521-5767 X-2455
Voicemail: 402-593-2455
Email: amy.lenhart@merck.com

Amber Liss-Erk
Office: 800-521-5767 X-2403
Voicemail: 402-593-2403
Email: amber.liss@merck.com